

## REVIEW

# Distribution, extent of inter-annual variability and diet of the bloom-forming jellyfish *Rhizostoma* in European waters

M.K.S. LILLEY, J.D.R. HOUGHTON AND G.C. HAYS

Institute of Environmental Sustainability, Swansea University, Singleton Park, Swansea, SA2 8PP, UK

*Jellyfish (Cnidaria: Scyphozoa) are increasingly thought to play a number of important ecosystem roles, but often fundamental knowledge of their distribution, seasonality and inter-annual variability is lacking. Bloom forming species, due to their high densities, can have particularly intense trophic and socio-economic impacts. In northern Europe it is known that one particularly large (up to 30 kg wet weight) bloom forming jellyfish is Rhizostoma spp. Given the potential importance, we set out to review all known records from peer-reviewed and broader public literature of the jellyfish R. octopus (Linnaeus) and R. pulmo (Macri) (Scyphozoa: Rhizostomae) across western Europe. These data revealed distinct hotspots where regular Rhizostoma spp. aggregations appeared to form, with other sites characterized by occasional abundances and a widespread distribution of infrequent observations. Surveys of known R. octopus hotspots around the Irish Sea also revealed marked inter-annual variation with particularly high abundances forming during 2003. The location of such consistent aggregations and inter-annual variances are discussed in relation to physical, climatic and dietary variations.*

**Keywords:** historical distributions, gelatinous zooplankton, aggregation, ecosystem shift, Mar Menor, *Rhysostoma*, *Rhisostoma*, *Rhyzostoma*

Submitted 18 January 2008; accepted 6 June 2008; first published online 9 September 2008

## INTRODUCTION

There is increasing evidence that a combination of climatic factors and human-induced change within the marine environment is increasing the prominence of jellyfish abundances from a number of locations worldwide (Mills, 2001; Hays *et al.*, 2005). For example, in the Bering Sea the biomass of jellyfish increased more than 10-fold during the 1990s (Brodeur *et al.*, 2002), although a subsequent reversal has occurred since 2000 despite further climatic warming (Purcell, 2005), whilst in the Benguela upwelling system jellyfish biomass (12.2 million tonnes) now exceeds the abundance of once-abundant fish (3.6 million tonnes) (Lynam *et al.*, 2006). Jellyfish populations may be increasing in biomass at existing sites, undergoing range expansions into new areas and in addition non-native invasive species may proliferate, with all these processes contributing to overall increases in

jellyfish prevalence (Mills, 2001). However, identifying these various scenarios is often hampered by inadequate information on spatial patterns of abundance and in particular the absence of historic data. Often old reports of jellyfish blooms are scattered in the literature and, as a result, the ecological importance of jellyfish within particular environments is often grossly underestimated (Mills, 2001).

Recent studies in the Irish Sea have used a variety of approaches to assess patterns of jellyfish abundance, including visual surveys from ships of opportunity (Doyle *et al.*, 2007), shoreline strandings (Houghton *et al.*, 2007) and aerial surveys (Houghton *et al.*, 2006a), (see also Purcell *et al.*, 2000; Graham *et al.*, 2003). These methods have revealed species-specific patterns of distribution (Doyle *et al.*, 2007). For example, within the Irish Sea the lions mane jellyfish (*Cyanea capillata*) appears to dominate cold northern waters, whilst the barrel jellyfish *Rhizostoma octopus* was found in massive numbers in large shallow embayments (Houghton *et al.*, 2006a, b; Doyle *et al.*, 2007). The large size of *Rhizostoma* (individuals up to 30 kg wet weight) and extent of the blooms identified in the Irish Sea (millions of individuals)

### Corresponding author:

M.K.S. Lilley

Email: 385124@swansea.ac.uk

suggest that this species has a large impact on the ecosystem, yet information on its wider distribution is fragmented. By drawing together data from peer-reviewed and broader public literature (dating back to 1838) we consider both temporal and spatial variations of the abundance of this species across western Europe and explore whether the high levels of abundance seen in selected sites in the Irish Sea, reflect a broader occurrence of hotspots. Further, we examine the first empirical evidence for the extent of inter-annual variability in the size of blooms and hence provide a starting point for assessing climate change impacts on this species. Finally we review the diet of this species and hence help to clarify its wider ecosystem role.

## MATERIALS AND METHODS

### Compiling historical reports of *Rhizostoma* spp. in European waters

We collated historical reports of *Rhizostoma* spp. Local and national newspaper archives (Table 1), wildlife trust records and personal accounts supplemented peer-reviewed records from the scientific literature. The nature of many of these reports could not enable definitive abundances to be estimated for each location, but it was possible to build up a picture of the known locations for large aggregations. *Rhizostoma octopus* (Linnaeus) and *R. pulmo* (Macri) (Scyphozoa: Rhizostomeae) are closely related and frequently misidentified, with the sole difference being the number of marginal lappets around the bell, generally 8 per octant in *R. pulmo* and 10 in *R. octopus*, but up to 16 have been recorded. A possible third species, *R. luteum*, has also been described (Russell, 1970) but no records were found during our investigations and the recognition of three species was questioned by Holst *et al.* (2007). Misidentification, renaming of species and genus only reports across the 180 year timescale were alleviated by pooling the two *Rhizostoma* species.

The recording frequency of species, such as *Rhizostoma* spp., is highly dependant on their effect on regional economies. The resulting data were treated at face-value as sightings, which we were unable to correct for effort, and may therefore have regional biases. Very few of the sightings, aside from recent peer-reviewed papers, should be regarded as providing any evidence for the absence of *Rhizostoma* spp.

### Assessing inter-annual variability using aerial survey data

Between 2003–2005 aerial surveys of jellyfish distribution were conducted for the entire Irish Sea (Houghton *et al.*, 2006a). Firstly, random transects were conducted across coastal and open-water areas, then once jellyfish aggregations had been identified, repeated line transects were employed over several months to assess the consistency of these features in space and time. Estimates of abundance were made using a simple visual count system with data expressed as number of individuals observed every 5 minutes. The five minute time frames translated to 7710 m<sup>2</sup> at a constant flying speed of 185 km hr<sup>-1</sup> and survey height of 152 m above sea level. Data were converted to abundances per 1000 m<sup>2</sup> with




Fig. 1. Reported occurrences of *Rhizostoma* spp. jellyfish in southern and western Europe from 1838–2007. Regions of rare sightings (white background), common sightings with occasional aggregations (grey), and regular aggregations (black) are highlighted.




Fig. 2. Mean *Rhizostoma octopus* densities (ind/1000 m<sup>2</sup>) and associated standard errors for the three 'hotspots' identified in Houghton *et al.* (2006a). Data for three years are shown for Carmarthen Bay and Tremadoc Bay and two years (2003–2004) for Rosslare harbour. Estimates are based only on survey effort within the three bays.

**Table 1.** Historical records of *Rhizostoma* spp. medusae from western Europe by region (Figure 1) (site numbers refer to symbols in Figure 1). Abundances are given where present. ‘This study’ refers to records extracted from outside the peer reviewed literature.

Site	Year	Location	Details of sightings
1	1980	Moray Firth	One 470 mm (Hay <i>et al.</i> , 1990)
2	1902	Isle of May	(Evans, 1916 in Russell, 1970)
	1913	North Berwick	(Evans, 1916 in Russell, 1970)
3	1960	Northumberland	(Fraser, 1961 in Russell, 1970)
4	Pre 1978	Norwegian coast	Regular sightings (Lid, 1979)
5	1873, 1877	Copenhagen	>1 (all Kramp, 1934)
	1896	Swedish coast	Large aggregations
	1898	Frederickshaven	A few medusae
	1916	Kristianiafjord	One
	1925	Jutland	Common this year
	1929	Frederickshaven	One 50 cm (very large) 20 November
	1933	Hantsholm/Hitshals	Common (to 40 cm) from early September until October
6	1925	Helgoland	Very few, (all Kramp, 1934 unless stated)
	1930	Helgoland	A few 17 September
	1933	Helgoland region	A few from late July, plenty off Elbe in mid-August, large numbers of big medusae during September
	1936–1938	Holland	Large numbers ~20 cm September/October, sometimes November too
	1964	Stuifdijk/Veerhaven	Intertidal nets catches July–October (Verwey, 1967)
	1992	Texel, Wadden Sea	23 October, largest stranding to date (December 2007) (Figure 4; S. Dijkse, personal communication)
	2003–2004	Helgoland	Individuals collected for research, no data on abundances (Holst & Jarms, 2006)
7	Pre 1954	Whitstable Bay, Thames estuary	Occasional in summer, to large size (Newell, 1954)
	1970	Isle of Grain, Kent	August—power station inlets blocked— <i>Rhizostoma</i> ? (this study)
	1976	Ostend, Belgium	June—large numbers (this study)
	2006	Nieuwpoortbad, Belgium	Images, no abundances (MarBEF, 2004)
8	1907	South coast	Large numbers (this study)
	1937	Hythe, Kent	Large numbers— <i>Rhizostoma</i> ? (this study)
	1987	South coast	July—some large medusae
	2002–2003	Cornwall to Shoreham, West Sussex	Large abundances (especially 2002) from March to October (this study)
9	1907	St Ives/Falmouth	Up to 900 mm (Valletin, 1907)
	1968	Plymouth	1 large one (Russell, 1970)
	1988	South-west region	Major <i>Rhizostoma</i> year (this study)
	1989	Mounts Bay	Large numbers (this study)
	1990	St Ives	April, fisherman’s nets full (this study)
	1993	South-west region	Major <i>Rhizostoma</i> year (this study)
	1999	Mounts Bay	Large numbers (this study)
	2002–2003	South-west coast	July—hot weather, plankton bloom and large numbers (this study)
	2006	Cornwall	Large numbers, see region 8 (this study) January, February and April—MCS aerial survey
10	1949–1965	Pembrokeshire	Seen 1949, 1950, 1952, 1955–1956, 1958, 1960 and 1965; abundant 1964 (Crothers, 1966)
	1960	Bideford	June—large numbers <i>Rhizostoma</i> ? (this study)
	1968	Minehead	Many stranded February/March (Russell, 1970)
10a	2003	Carmarthen Bay	Thousands August/September (Houghton <i>et al.</i> , 2006a)
	2004–2005		Fewer than 2003 (Houghton <i>et al.</i> , 2006a)
11	1966	Anglesey	February and May 1966 (Russell, 1970)
	2001	Tremadog Bay	Large numbers, hot weather (this study)
	2003–2005		Abundant (Houghton <i>et al.</i> , 2006a)
12	1894	Isle of Man	580 mm in April (Browne, 1895)
	1966	Irish Sea	>100 in 1/2 hour trawl (Russell, 1970)
12a		Solway Firth	July, 16–35 mm specimens caught for examination (Russell, 1970)
	1999	Lancashire	One stranded (McMillan, 1999)
	2004	Solway Firth	44 trawled in intensive beam trawl survey (Axelsson <i>et al.</i> , 2006)
	2005		Abundant (Houghton <i>et al.</i> , 2006b)
13	1901	Cumrae, Clyde	2 seen September/October (Browne, 1905)
	1992	Hunterston, Clyde	Power station inlets blocked (Houghton <i>et al.</i> , 2006a)
14	1838	Belfast Bay	(Thompson, 1840 in Boyd <i>et al.</i> , 1973) autumn (O’Connor & McGrath, 1978)
	1857	Dublin	Greene 1857 in Boyd <i>et al.</i> , 1973
	1865		100s, March (Baily, 1865 in Boyd <i>et al.</i> , 1973)
	1886	Rush, Dublin	(Haddon, 1886 in Boyd <i>et al.</i> , 1973)
	1906	Lough Foyle	Two seen, February (O’Connor & McGrath, 1978)

Continued

Table 1. Continued

Site	Year	Location	Details of sightings
	1909	Strangford Lough	(Colgan, 1914 and Allen, 1936 in O'Connor & McGrath, 1978)
	1936–1937	Rathlin Island, County Antrim	Frequent (McDonald & McMillan, 1951 in Williams, 1954)
	Pre 1954	Strangford Lough	Regular (Williams, 1954); June–October (O'Connor & McGrath, 1978)
		Laytown, County Louth	Summer–mid-October (O'Connor & McGrath, 1978)
	2006	Off Strangford Lough	November, offshore, 4 adults (72, 40, 35 and 35 cm) trawled from bottom (this study)
	2007	Off Howth, Dublin	February, offshore, two 80 cm medusae trawled (this study) North Channel, February, 50–60 cm medusa trawled (this study)
14a	1976	Rosslare harbour	February, March and October (O'Connor & McGrath, 1978)
	2003–2004		Abundant (Houghton <i>et al.</i> , 2006a)
15	1838	Youghal, Cork	Autumn (Thompson, 1856 in Boyd <i>et al.</i> , 1973)
	1896	Valencia	1 in October (Browne, 1896 in O'Connor & McGrath, 1978)
	1900		Many, September (Browne, 1900 in Boyd <i>et al.</i> , 1973)
	1905		August–October (Delap & Delap, 1906 in Boyd <i>et al.</i> , 1973)
	1924		Many, September/October (Boyd <i>et al.</i> , 1973)
	1976		Mid-August–mid-September (O'Connor & McGrath, 1978)
		Dungaven, County Waterford	July–September (O'Connor & McGrath, 1978)
		Courtmacsherry Bay County Cork	August (O'Connor & McGrath, 1978)
		Kinsale, County Cork	(O'Connor & McGrath, 1978)
16	1844	West coast	(Thompson, 1856 in Boyd <i>et al.</i> , 1973)
	Pre 1972	Galway	July–September, may be common (Boyd <i>et al.</i> , 1973)
	1976	Dingle Bay, County Kerry	April–May (O'Connor & McGrath, 1978)
		Liscannor, County Clare	May–mid-November (O'Connor & McGrath, 1978)
		Galway Bay	May–October (O'Connor & McGrath, 1978)
		Killary harbour, County Galway	May (O'Connor & McGrath, 1978)
		Achill harbour, County Mayo	Summer (O'Connor & McGrath, 1978)
		Killala Bay, North Mayo	Summer (O'Connor & McGrath, 1978)
		Donegal Bay	June–September (O'Connor & McGrath, 1978)
17	1896	Trevignon	One found in turtle stomach (Duron, 1978)
	1977–1978	La Rochelle	Abundant, especially 1978—10–30 cm, 1.5 miles diameter aggregation, August–September (Duron, 1978)
	1982		July, one 20 kg, 87 cm (this study)
	1996		July—large numbers (this study)
	1998		July—large numbers (this study)
	2003		August—exceptional number in the hot weather (this study)
18	1982	Hendaye, Bilbao	December, 20 tonnes in fishing trawl (this study)
19	1993–2001	Mar Menor ( <i>Rhizostoma pulmo</i> )	Big aggregations <40 cm annually May–July (Mas, 1999; Perez-Ruzafa <i>et al.</i> , 2002)
	2002–2004		No direct data
	2005		Fewer than normal—1993–2004 average (this study)
	2006		Large numbers 35–40 cm (this study)
		Near Barcelona	>1500 stranded (this study)
20	1898–1914	Ligurian Sea	Regular population recorded particularly from October–January, no aggregations (Morand & Dallot, 1985)
	2003	Tuscany	July—>100,000 stranded (this study)

standard errors calculated from these. Four key hotspots for *Rhizostoma octopus* were identified: (1) Carmarthen Bay (number of 1000 m<sup>2</sup> survey points per year 2003–05 respectively = 83, 63 and 22); (2) Tremadoc Bay (N = 31, 59 and 48); (3) Rosslare harbour (N = 28, 51 and 0); and (4) Solway Firth (Figure 1; locations 10a, 11, 14a and 12a respectively); although data for the site 4 was only collected during a single year (Houghton *et al.*, 2006a, b). Two-sample *t*-tests compared the inter-annual variance in abundance of *R. octopus* over three consecutive years for each of sites 1–3.

### Prey items for *Rhizostoma* spp

We considered gut content data from a range of previous studies. Prey size and dietary percentage was considered for five scyphozoan species: *Cotylorhiza tuberculata* and

*Rhizostoma pulmo* (data obtained from Perez-Ruzafa *et al.*, 2002); *Chrysaora quinquecirrha* (Purcell, 1992); *Aurelia aurita* (Hamner *et al.*, 1982; Sullivan *et al.*, 1994; Graham & Kroutil, 2001; Ishii & Tanaka, 2001; Barz & Hirche, 2005) and *Cyanea capillata* (Fancett, 1988; Brewer, 1989; Purcell, 2003). Dietary compositions in each study were predominantly recorded as prey abundance consumed per medusa. These values were converted to proportion of diet for each study and then averaged by species where more than one study was referred to. Prey species were commonly identified by name rather than size-class requiring establishment of size-ranges from published sources (Table 2). In the absence of definitive prey sizes, data were classified into 200 µm prey length bins up to 1000 µm to reduce errors, with larger bins for species greater than 1000 µm.

**Table 2.** Length-class of scyphozoan prey items (Figure 3), recorded length and published source.

Length-class ( $\mu\text{m}$ )	Species	Recorded length ( $\mu\text{m}$ )	Source
0–200	Tintinnids	<100	(Capriulo & Carpenter, 1983)
	Coscinodiscus	100–150	(Alpine & Cloern, 1985)
	Bivalve veligers	100–260	(Purcell <i>et al.</i> , 1991)
	Rotifers	<200	(Stemberger & Gilbert, 1985)
	Copepod nauplii	150	(Munk & Kiorboe, 1985)
200–400	Larvacean (trunk)	200–500	(Uye & Ichino, 1995)
400–600	Cladocerans	<500	(Zaret & Kerfoot, 1975)
600–800	Harpacticoids	400–900	(Sun & Fleeger, 1995)
800–1000	Copepods	<1000	(Graham & Kroutil, 2001)
1000–2000	Copepods	>1000	(Graham & Kroutil, 2001)
	Crab zoea	1000–2000	(Gore, 1968)
>2000	Fish eggs	1000–2000	(Pauly & Pullin, 1988)
	Hydromedusae	>6000–13000	(Nicholas & Frid, 1999)

## RESULTS

### Historical reports of *Rhizostoma* spp. in European waters

Spatial and temporal records of *Rhizostoma* spp. reveal both irregular sightings of individual medusae and reports of large aggregations or blooms (Figure 1). Treating these two groupings separately and chronologically, individual reports from the 19th and early 20th Centuries suggest a broad, but sporadic, distribution across European waters. Pre-1900 *Rhizostoma* had been reported as widely as Copenhagen, the Irish Sea, around the coast of west, south and eastern Ireland and in the Bay of Biscay (see Russell, 1970 for examples). The sole Mediterranean Sea recording, from the Ligurian Sea off north-western Italy, was reported annually particularly in autumn and winter from 1898 until 1914 (Morand & Dallot, 1985). Apart from this Ligurian population, the early 1900s were fairly sparsely populated with reports of *Rhizostoma*. Notably this period included the first couple of reports from eastern Scotland and one from western Scotland, but the reduction in frequency of records in the formerly richly recorded Irish waters may represent an overall reduction in the production of *Rhizostoma* spp. or may simply reflect a reduced sighting effort during this period. The 1930s saw a resurgence of reports, predominantly in the Helgoland and Kattegat regions (Kramp, 1934) with frequent observations from south-west Ireland in 1937–1938 (McDonald & McMillan, 1951 in Williams, 1954). A complete absence of reports during the Second World War was followed by the first report from the Thames estuary and almost annual observations from the field station at Dale Fort in Pembrokeshire (Crothers, 1966). The 1960s and 1970s saw a series of observations from Ireland, a few from south-west England, the only report from Norway and the first since 1896 in the Bay of Biscay (see Table 1). Reports obtained for the last two decades of the 20th Century were predominantly focused on southern England and the Bay of Biscay with many reports of blooms rather than individuals. This pattern has continued into the 21st Century with a considerable number of reports already in the first seven years, the furthest north of these within the Irish Sea (Houghton *et al.*, 2006b).

Taking European waters as a whole we categorized the frequency of years with large aggregations in the reporting of *Rhizostoma*. Table 1 summarizes these findings. Chronologically, prior to the First World War there had

only been mass aggregations reported from Dublin in 1865, off the Swedish coast in 1896, south-west Ireland in 1900, and the south coast of England in 1907. Inter-war, south-west Ireland experienced a second aggregation in 1924 and the Helgoland/Jutland region of the North Sea saw blooms in 1925 and from 1936–1938, with the 1937 aggregation also spotted on the English coast. An absence of reported blooms during the 1940s and 1950s was broken in 1960 and 1968 by strandings on the south-west England peninsular and a large haul of *Rhizostoma* fished from the Irish Sea during 1966 (Russell, 1970). Two records exist of European *Rhizostoma* blocking power stations cooling water inlets—on the Isle of Grain, Thames estuary, in 1970 and Hunterston, south-west Scotland, in 1992—illustrating the magnitude of the associated blooms and the bulk of the medusae. A report of a bloom from Belgium in 1976 was more than matched by a huge abundance that was reported spreading clockwise around the Irish coasts (O'Connor & McGrath, 1978). The following year aggregations were reported from the Bay of Biscay for the first time with further reports in 1978 and 1982 (twenty tonnes caught by a fisherman), but there was an absence of blooms further north during this period. Since the late 1980s frequent clustered reports from the English Channel have been interspersed with Bay of Biscay reports in 1996, 1998 and 2003. Of these only that in 2003 appears to have been mimicked across northern Europe with three Irish Sea embayments (Carmarthen, Tremadog and Rosslare (Houghton *et al.*, 2006a, b)), the English Channel and the Bay of Biscay involved. In the Mediterranean, the lagoon habitat of Mar Menor saw annual blooms from 1993–2001 and again in 2006, with open sea blooms reported in 2003 (Tuscany) and 2006 (Barcelona). These reports would appear to suggest that the frequency of bloom formations has increased, especially in the late 20th Century, however an obvious absence of data during the World Wars gives some indication of the sporadic nature of sighting reports.

### Assessing inter-annual variability using aerial survey data

Aerial surveys across the Irish Sea during 2003–2005 revealed four hotspot locations for *Rhizostoma* spp (Houghton *et al.*, 2006a): (1) Carmarthen Bay; (2) Tremadoc Bay; (3) the Solway Firth; and (4) Rosslare harbour (Figure 1; 10a, 11,

12a and 14a). Site 3 was only identified during 2005 and was therefore excluded from inter-annual estimates.

Kruskal–Wallis analysis revealed significant higher abundances during 2003 compared to other years ( $H_{1,2} = 19.32$ ,  $P < 0.001$  data adjusted for ties). Mann–Whitney  $U$ -tests revealed significant reductions in abundance of *Rhizostoma* between 2003–2004 and 2003–2005 (Figure 2) ( $P < 0.001$  in each case) while no difference was found overall between 2004 and 2005 ( $P > 0.05$ ). Within sites Kruskal–Wallis tests confirmed difference between years for Site 1 ( $H_{1,2} = 7.03$ ,  $P = 0.03$ ) and Site 2 ( $H_{1,2} = 17.63$ ,  $P < 0.001$ ), but not Site 4 ( $H_{1,2} = 3.13$ ,  $P = 0.07$  all results adjusted for ties). Pair-wise Mann–Whitney  $U$ -test comparisons at Site 1 found a significant reduction between 2003 and 2004 ( $P = 0.011$ ) but no difference between 2005 and either year. Pair-wise comparisons at Site 2 found significant reductions from 2003 to 2004 and 2005 ( $P = 0.002$  and  $0.001$  respectively) but not between the later years. In short there were significant interannual variations in *R. octopus* abundances with higher numbers in 2003 compared to other years.

### Prey items for *Rhizostoma* spp

The two species present in the coastal lagoon of Mar Menor in Región de Murcia, Spain, *R. pulmo* and *Cotylorhiza tuberculata*, consume distinctly different size prey compared to other commonly studied scyphozoan species. In Mar Menor the predominant species preyed upon were the diatoms *Asterionella* and *Coscinodiscus* (Perez-Ruzafa *et al.*, 2002),  $< 200 \mu\text{m}$  size-class (Figure 3), with tintinnids and veligers also forming significant components. Copepods made up 0.5% and 3.17% of the diets respectively. *Aurelia aurita*, *C. quinquecirrha* and *C. capillata* were all found to consume a diverse mix of larvaceans (200–400  $\mu\text{m}$  trunk length), copepods (800–1000  $\mu\text{m}$  and 1000–2000  $\mu\text{m}$ ), cladocerans species such as *Bosmina* (400–600  $\mu\text{m}$ ) and fish eggs (1000–2000  $\mu\text{m}$ ) among other prey.

## DISCUSSION

### Compiling historical reports of *Rhizostoma* spp. in European waters

Our assessment suggests that blooms of *Rhizostoma* are widespread across Europe. Clearly our work draws heavily on the use of data from the grey literature, which is necessitated due to absence of large scale systematic surveys of jellyfish abundance across the area of interest. The use of the grey literature can be defended here since *Rhizostoma* is an easy to identify genus, and due to its large size the sightings of individuals might often be a newsworthy observation. However, despite this the recording frequency will depend heavily on the effect of the medusae on regional economies, in particular fisheries and tourism, with consequential variation between seasons, regions and popularity of locations for observers. Clearly not all occurrences of *Rhizostoma* will be reported and so this material provides essentially information on the minimum possible occurrence of bloom locations. Similarly while the large aggregations around the Helgoland region of The Netherlands spilling into the Kattegat in 1933–1934 (Kramp, 1934) and encircling Ireland during 1976




Fig. 3. Prey length and dietary percentage for five scyphozoan jellyfish extracted from published gut contents studies. Data for *Aurelia aurita* (a), *Cyanea capillata* (b) and *Chrysaora quinquecirrha* (c) were averaged from published data. *Cotylorhiza tuberculata* (d) and *Rhizostoma pulmo* (e) data were reproduced from Perez-Ruzafa *et al.* (2002); *Chrysaora quinquecirrha* data were averaged from populations described in Purcell (1992); *Aurelia aurita* data were averaged from Sullivan *et al.* (1994), Graham & Kroutil (2001), Ishii & Tanaka (2001), Barz & Hirche (2005) and Hamner *et al.* (1982); and *Cyanea capillata* data from Brewer (1989), Fancett (1988) and Purcell (2003).

(O'Connor & McGrath, 1978) were reported in the peer-reviewed literature, the scarce reports from these areas in other years, might simply reflect a lack of scientific study rather than an absence of the jellyfish themselves.

We have previously used aerial surveys to show that bloom locations for *Rhizostoma* are consistent from one year to



**Fig. 4.** A mass stranding of *Rhizostoma octopus* near Paal 12, Texel, Dutch Wadden Sea on 23 October 1992. This large stranding went otherwise unrecorded (Sytske Dijkse/Foto Fitis).

another (Houghton *et al.*, 2006b). Based on this regional evidence we would suggest that sites across Europe where occasional blooms have been reported, most probably reflect hotspots for blooms, to varying degrees, every year but with only sporadic reporting of these events (Figure 4). This suggestion could clearly be tested by focused surveys at those sites (e.g. the north-western Mediterranean and off the Elbe estuary at Helgoland, Holland) which would be straightforward to perform using established methodologies (Purcell *et al.*, 2000; Graham *et al.*, 2003; Houghton *et al.*, 2006a).

Our compilation of historic records show that the recent blooms of *Rhizostoma* recorded in the Irish Sea reflect a broader occurrence of blooms, with hotspots also being found around La Rochelle in the Bay of Biscay, and Mar Menor, Spain. The particular conditions within these areas that are conducive to blooms of *Rhizostoma* are still not clear, but all these areas are semi-enclosed bay areas that receive appreciable freshwater and nutrient input from rivers. In the case of Mar Menor, the freshwater regime significantly increased post-dredging of the entrances, but is still low compared to other Mediterranean lagoons, suggesting controls such as nutrient loads may have more of an effect than hydrographic or trophic variables (Pérez-Ruzafa *et al.*, 2005). Further analysis is clearly needed to pin-point the underlying reasons for why blooms occur in these areas and such work might also point the way to where other, as yet unreported blooms of *Rhizostoma* are to be expected.

While targeted studies have shown the *Rhizostoma* spp. hotspots may be consistent year-after-year, there is clearly a large degree of variability in the extent of blooms. For example, across our aerial survey data there were

approximately three and nine times more *R. octopus* in 2003 than 2004 and 2005 respectively. However, Carmarthen Bay alone had a 478 times difference between 2003 and 2005. The processes driving this inter-annual variability are obscure. However, inter-annual variability in the abundance of jellyfish has been noted elsewhere. For example Purcell *et al.* (2000) showed that the abundance of *Aurelia labiata* in Alaska varied by two-fold between years. Similarly a 17-year times-series of jellyfish abundance for the northern North Sea derived from fish trawl data showed that inter-annual variability in the abundance of *Aurelia aurita*, *Cyanea capillata* and *C. lamarckii* was correlated to the winter North Atlantic Oscillation Index (NAOI) (Lynam *et al.*, 2004, 2005). Our aerial records of *R. octopus* revealed a matching pattern of reduced abundance between 2003 and 2005 in bays over 100 km apart. Furthermore there were reports of *Rhizostoma* spp. aggregations and strandings across Europe in the public press during 2003. *Rhizostoma pulmo* was stranded in hundreds of thousands on the shore of Tuscany, Italy (10 July 2003, Neue Zuercher Zeitung, Zurich) and 'exceptional numbers' in western France near La Rochelle (15 August 2003, Sud Ouest, Bordeaux) in addition to the annual abundances in Mar Menor. The implication is that broad-scale processes were affecting levels of abundance rather than site-specific processes. This finding is consistent with the proposed broad-scale changes in jellyfish and other zooplankton abundance reported by Moline *et al.* (2004), Hay (2006) and Lynam *et al.* (2006). Of interest is the fact that across Europe, 2003 was an exceptionally hot year with temperatures attaining levels predicted by climatic models for the late 21st Century (Beniston, 2004;

Luterbacher *et al.*, 2004; Schär & Jendritzky, 2004; Ciais *et al.*, 2005) with widespread implications for species distributions (e.g. Battisti *et al.*, 2006; Jiguet *et al.*, 2006; Mouthon & Daufresne, 2006). With high abundances of *Rhizostoma* spp. correlating with the high temperatures in Europe we speculate that warmer summers will lead to increasing *Rhizostoma* blooms.

The lifecycle of *R. octopus* was published by Holst *et al.* (2007) from cultures maintained within the laboratory providing, for the first time, clear images of the 2 mm size of the polyps. The authors specified a decrease in temperature from 15°C to 10°C or rises from 5–10°C or 10–15°C was required to stimulate strobilation of the polyps. For *R. octopus* these temperatures would provide a reproductive range across northern Europe. However, for the predominantly southern European species *R. pulmo*, if water temperature is the key stimuli to strobilation as hypothesized by Pérez-Ruzafa (1997 cited in Kingsford *et al.*, 2000) the temperatures at which strobilation is stimulated are likely to be higher. A further study utilized ephyrae to settle on the undersides of substrates and a preference for plastic compared to other natural and anthropogenic materials tested (Holst & Jarms, 2006). However, neither of these studies identified locations or typical substrates where the polyp stages could be found in the natural environment. This lack of information on the polyp stage of free-living individuals remains a key stumbling block to explain spatio-temporal patterns of occurrence for *Rhizostoma*.

### Prey items for *Rhizostoma* spp

Gelatinous zooplankton aggregations form predominantly mono-species aggregations (Mills, 2001) resulting from the variability of prey and the conditions required for population increases. Most species of jellyfish employ banks of nematocysts in tentacles for capturing planktonic prey. The scyphozoan *Rhizostoma* spp., by contrast, does not possess long trailing tentacles, but filters prey through a mass of eight oral arms hanging below the bell. Perez-Ruzafa *et al.* (2002) provided gut contents measurements for *R. pulmo* in Mar Menor, Spain. Their data would indicate a predominantly small-size phytoplanktonic diet, contrasting with the zooplankton dominated diet of *A. aurita*, *C. quinquecirrha* and *C. capillata* (Figure 3). The jellyfish abundance of Mar Menor correlated well with increased diatom abundances and nitrate levels (Perez-Ruzafa *et al.*, 2002) from 1993–2006 (few *R. pulmo* were present during 2005; Table 1).

Perez Ruzafa *et al.* (2002) also highlighted the regular location of *R. pulmo* aggregations in the south-western part of the lagoon closest to the freshwater (and nutrient) inflows. The phytoplankton standing stock, varied by season, was particularly dominated by diatoms (*Cyclotella*, *Chaetoceros* and *Nitzschia* spp.) between spring and autumn, thereby apparently forming a large part of the medusa's diet. It is this standing stock caused by the regular nutrient input and warm shallow waters (Gilbert, 2001) that is most likely to be the driving factor behind the regular abundances of *Rhizostoma* in Mar Menor. The dietary consumption of species is important for calculating the impact of gelatinous zooplankton on the ecosystem as a whole. Studies have suggested that jellyfish may be an important conduit for energy flow through marine systems

(Lucas *et al.*, 1997). Ultimately the ecosystem role of jellyfish may be better understood by parameterizing jellyfish within ecosystem models. This parameterization will require information on levels of abundance but also jellyfish prey and rate processes such as ingestion.

### ACKNOWLEDGEMENTS

M.K.S.L. was supported by a studentship from the Natural Environment Research Council of the UK (NERC). J.D.R.H. and G.C.H. were supported by a NERC Urgency grant. Thanks also to two anonymous referees for their constructive criticism in improving the final manuscript.

### REFERENCES

- Alpine A.E. and Cloern J.E. (1985) Differences in *in vivo* fluorescence yield between three phytoplankton size-classes. *Journal of Plankton Research* 7, 381–390.
- Axelsson M., Dewey S., Tourell A. and Karpouzli E. (2006) *Site condition monitoring—the sublittoral sandbanks of the Solway Firth*. Scottish Natural Heritage Commissioned Report, No. 155 (ROAME No. F02AA409), pp. 111.
- Barz K. and Hirche H.J. (2005) Seasonal development of scyphozoan medusae and the predatory impact of *Aurelia aurita* on the zooplankton community in the Bornholm Basin (central Baltic Sea). *Marine Biology* 147, 465–476.
- Battisti A., Stastny M., Buffo E. and Larsson S. (2006) A rapid altitudinal range expansion in the pine processionary moth produced by the 2003 climatic anomaly. *Global Change Biology* 12, 662–671.
- Beniston M. (2004) The 2003 heat wave in Europe: a shape of things to come? An analysis based on Swiss climatological data and model simulations. *Geophysical Research Letters* 31.
- Boyd R.J., O'Ceidigh P. and Wilkinson A. (1973) Investigations of the plankton of the west coast of Ireland—VI. Pelagic Cnidaria of the Galway Bay area 1956–72, with revision of previous records for these species in Irish inshore waters. *Proceedings of the Royal Irish Academy* 73B, 383–403.
- Brewer R.H. (1989) The annual pattern of feeding, growth, and sexual reproduction in *Cyanea* (Cnidaria: Scyphozoa) in the Niantic River Estuary, Connecticut. *Biological Bulletin. Marine Biological Laboratory, Woods Hole* 176, 272–281.
- Brodeur R.D., Sugisaki H. and Hunt G.L. (2002) Increases in jellyfish biomass in the Bering Sea: implications for the ecosystem. *Marine Ecology Progress Series* 233, 89–103.
- Browne E.T. (1895) Report on the medusae of the LMBC District (Isle of Man). *Transactions of the Liverpool Biological Society* 9, 243–286.
- Browne E.T. (1905) A report on the medusae found in the Firth of Clyde (1901–2). *Proceedings of the Royal Society of Edinburgh* 25, 736–78.
- Capriulo G.M. and Carpenter E.J. (1983) Abundance, species composition and feeding impact of tintinnid micro-zooplankton in central Long Island Sound. *Marine Ecology Progress Series* 10, 277–288.
- Ciais P., Reichstein M., Viovy N., Granier A., Ogee J., Allard V., Aubinet M., Buchmann N., Bernhofer C. and Carrara A. (2005) Europe-wide reduction in primary productivity caused by the heat and drought in 2003. *Nature* 437, 529–533.
- Crothers J.H. (ed.) (1966) *Dale Fort marine fauna*, 2nd edition. London: Field Studies Council.

- Doyle T.K., Houghton J.D.R., Buckley S.M., Hays G.C. and Davenport J. (2007) The distribution of five scyphozoan jellyfish species across a temperate coastal environment. *Hydrobiologia* 579, 29–39.
- Duron M. (1978) *Contribution à l'étude de la biologie de Dermochelys coriacea (Linné) dans les Pertuis Charentais*. Bordeaux: Université de Bordeaux.
- Fancett M.S. (1988) Diet and prey selectivity of scyphomedusae from Port Phillip Bay, Australia. *Marine Biology* 98, 503.
- Gilbert J. (2001) Seasonal plankton dynamics in a Mediterranean hypersaline coastal lagoon: the Mar Menor. *Journal of Plankton Research* 23, 207–218.
- Gore R.H. (1968) The larval development of the commensal crab *Polyonyx gibbesi* Haig, 1956 (Crustacea: Decapoda). *Biological Bulletin. Marine Biological Laboratory, Woods Hole* 135, 111–129.
- Graham W.M. and Kroulitz R.M. (2001) Size-based prey selectivity and dietary shifts in the jellyfish, *Aurelia aurita*. *Journal of Plankton Research* 23, 67–74.
- Graham W.M., Martin D.L., Felder D.L., Asper V.L. and Perry H.M. (2003) Ecological and economic implications of a tropical jellyfish invader in the Gulf of Mexico. *Biological Invasions* 5, 53–69.
- Hamner W.M., Gilmer R.W. and Hamner P.P. (1982) The physical, chemical, and biological characteristics of a stratified, saline, sulfide lake in Palau. *Limnology and Oceanography* 27, 896–909.
- Hay S. (2006) Marine ecology: gelatinous bells may ring change in marine ecosystems. *Current Biology* 16, R679–R682.
- Hay S.J., Hislop J.R.G. and Shanks A.M. (1990) North Sea Scyphomedusae—summer distribution, estimated biomass and significance particularly for O-group gadoid fish. *Netherlands Journal of Sea Research* 25, 113–130.
- Hays G.C., Richardson A.J. and Robinson C. (2005) Climate change and marine plankton. *Trends in Ecology and Evolution* 20, 337.
- Holst S. and Jarms G. (2006) Substrate choice and settlement preferences of planula larvae of five Scyphozoa (Cnidaria) from German Bight, North Sea. *Marine Biology* 151, 863–871.
- Holst S., Sötje I., Tiemann H. and Jarms G. (2007) Life cycle of the rhizostome jellyfish *Rhizostoma octopus* (L.) (Scyphozoa, Rhizostomeae), with studies on cnidocysts and statoliths. *Marine Biology* 151, 1695–1710.
- Houghton J.D.R., Doyle T.K., Davenport J. and Hays G.C. (2006a) Developing a simple, rapid method for identifying and monitoring jellyfish aggregations from the air. *Marine Ecology Progress Series* 314, 159–170.
- Houghton J.D.R., Doyle T.K., Wilson M.W., Davenport J. and Hays G.C. (2006b) Jellyfish aggregations and leatherback turtle foraging patterns in a temperate coastal environment. *Ecology* 87, 1967–1972.
- Houghton J.D.R., Doyle T.K., Lilley M.K.S., Wilson R.P., Davenport J. and Hays G.C. (2007) Stranding events provide indirect insights into seasonality and persistence of jellyfish (Cnidaria: Scyphozoa) medusae. *Hydrobiologia* 589, 1–13.
- Ishii H. and Tanaka F. (2001) Food and feeding of *Aurelia aurita* in Tokyo Bay with an analysis of stomach contents and a measurement of digestion times. *Hydrobiologia* 451, 311–320.
- Jiguet F., Julliard R., Thomas C.D., Dehorter O., Newson S.E. and Couvet D. (2006) Thermal range predicts bird population resilience to extreme high temperatures. *Ecology Letters* 9, 1321–1330.
- Kingsford M.J., Pitt K.A. and Gillanders B.M. (2000) Management of jellyfish fisheries, with special reference to the Order Rhizostomeae. *Oceanography and Marine Biology* 38, 85–156.
- Kramp P.L. (1934) An exceptional occurrence of *Rhizostoma octopus* and *Chrysaora hysoscella* in the Danish waters in 1933. *Journal du Conseil* 9, 211–221.
- Lid G. (1979) The occurrence of jellyfish (Scyphozoa) along the Norwegian coast. *Fauna (Blindern)* 32, 129–136.
- Lucas C.H., Hirst A.G. and Williams J.A. (1997) Plankton dynamics and *Aurelia aurita* production in two contrasting ecosystems: comparisons and consequences. *Estuarine, Coastal and Shelf Science* 45, 209–219.
- Luterbacher J., Dietrich D., Xoplaki E., Grosjean M. and Wanner H. (2004) European seasonal and annual temperature variability, trends, and extremes since 1500. *Science* 303, 1499–1503.
- Lynam C.P., Hay S.J. and Brierley A.S. (2004) Interannual variability in abundance of North Sea jellyfish and links to the North Atlantic Oscillation. *Limnology and Oceanography* 49, 637–643.
- Lynam C.P., Hay S.J. and Brierley A.S. (2005) Jellyfish abundance and climatic variation: contrasting responses in oceanographically distinct regions of the North Sea, and possible implications for fisheries. *Journal of the Marine Biological Association of the United Kingdom* 85, 435–450.
- Lynam C.P., Gibbons M.J., Axelsen B.E., Sparks C.A.J., Coetzee J., Heywood B.G. and Brierley A.S. (2006) Jellyfish overtake fish in a heavily fished ecosystem. *Current Biology* 16, 492–493.
- MarBEF (2004) *European Register of Marine Species*. Available online at <http://www.marbef.org/data/erms.php>. [Accessed 5 December 2007.]
- Mas J. (1999) *Estudio de la dinamica de poblaciones de las medusas Cetylorthiza tuberculata, Rhizostoma pulmo y Aurelia aurita en el mar Menor y de su problematica asociada*. Instituto Espanol de Oceanografia, internal report, pp. 101.
- McMillan N. (1999) Actinians (sea-anemones) beached in large numbers on the south Lancashire coast. *Naturalist* 124, 55–56.
- Mills C.E. (2001) Jellyfish blooms: are populations increasing globally in response to changing ocean conditions? *Hydrobiologia* 451, 55–68.
- Moline M.A., Claustre H., Frazer T.K., Schofield O. and Vernet M. (2004) Alteration of the food web along the Antarctic Peninsula in response to a regional warming trend. *Global Change Biology* 10, 1973–1980.
- Morand P. and Dallot S. (1985) Variations annuelles et pluriannuelles de quelques especes du macroplankton cotier de la mer Ligure (1898–1914). *Rapports et Proces-Verbaux des Réunions. Commission Internationale Pour l'Exploration Scientifique de la Mer Méditerranée* 29, 285–287.
- Mouthon J. and Daufresne M. (2006) Effects of the 2003 heatwave and climatic warming on mollusc communities of the Saone: a large lowland river and of its two main tributaries (France). *Global Change Biology* 12, 441.
- Munk P. and Kiorboe T. (1985) Feeding behaviour and swimming activity of larval herring (*Clupea harengus*) in relation to density of copepod nauplii. *Marine Ecology Progress Series* 24, 15–21.
- Newell G.E. (1954) The marine fauna of Whitstable. *Annals and Magazine of Natural History* 7, 321–350.
- Nicholas K.R. and Frid C.L.J. (1999) Occurrence of hydromedusae in the plankton off Northumberland (western central North Sea) and the role of planktonic predators. *Journal of the Marine Biological Association of the United Kingdom* 79, 979–992.
- O'Connor B. and McGrath D. (1978) On the occurrence of the scyphozoan *Rhizostoma octopus* (L.) around the Irish coast in 1976. *Irish Naturalists Journal* 19, 261–263.
- Pauly D. and Pullin R.S.V. (1988) Hatching time in spherical, pelagic, marine fish eggs in response to temperature and egg size. *Environmental Biology of Fishes* 22, 261–271.

- Perez-Ruzafa A.** (1997) *Estudio de la dinámica de las poblaciones de medusas en el Mar Menor, problemática asociada y búsqueda de soluciones* Final report of the Investigation Ecología y Ordenación de Ecosistemas Marinos Costeros Group. Department of Ecology and Hydrology, University of Murcia, Spain, pp. 234.
- Perez-Ruzafa A., Gilabert J., Gutierrez J.M., Fernández A.I., Marcos C. and Sabah S.** (2002) Evidence of a planktonic food web response to changes in nutrient input dynamics in the Mar Menor coastal lagoon, Spain. *Hydrobiologia* 475, 359–369.
- Pérez-Ruzafa A., Fernández A.I., Marcos C., Quispe J.I. and García-Charton J.A.** (2005) Spatial and temporal variations of hydrological conditions, nutrients and chlorophyll *a* in a Mediterranean coastal lagoon (Mar Menor, Spain). *Hydrobiologia* 550, 11–27.
- Purcell J.E.** (1992) Effects of predation by the scyphomedusan *Chrysaora quinquecirrha* on zooplankton populations in Chesapeake Bay, USA. *Marine Ecology Progress Series* 87, 65–76.
- Purcell J.E.** (2003) Predation on zooplankton by large jellyfish, *Aurelia labiata*, *Cyanea capillata* and *Aequorea aequorea*, in Prince William Sound, Alaska. *Marine Ecology Progress Series* 246, 137–152.
- Purcell J.E.** (2005) Climate effects on formation of jellyfish and ctenophore blooms: a review. *Journal of the Marine Biological Association of the United Kingdom* 85, 461–476.
- Purcell J.E., Cresswell F.P., Cargo D.G. and Kennedy V.S.** (1991) Differential ingestion and digestion of bivalve larvae by the scyphozoan *Chrysaora quinquecirrha* and the ctenophore *Mnemiopsis leidyi*. *Biological Bulletin. Marine Biological Laboratory, Woods Hole* 180, 103–111.
- Purcell J.E., Brown E.D., Stokesbury K.D.E., Halderson L.H. and Shirley T.C.** (2000) Aggregations of the jellyfish *Aurelia labiata*: abundance, distribution, association with age-0 walleye pollock, and behaviours promoting aggregation in Prince William Sound, Alaska, USA. *Marine Ecology Progress Series* 195, 145–158.
- Russell F.S.** (1970) *The Medusae of the British Isles. Volume II—Pelagic Scyphozoa*. Cambridge: Cambridge University Press.
- Schär C. and Jendritzky G.** (2004) Climate change: hot news from summer 2003. *Nature* 432, 559–560.
- Stemberger R.S. and Gilbert J.J.** (1985) Body size, food concentration, and population growth in planktonic rotifers. *Ecology* 66, 1151–1159.
- Sullivan B.K., Garcia J.R. and Klein-MacPhee G.** (1994) Prey selection by the scyphomedusan predator *Aurelia aurita*. *Marine Biology* 121, 335–341.
- Sun B. and Fleeger J.W.** (1995) Sustained mass culture of *Amphiascoides atopus* a marine harpacticoid copepod in a recirculating system. *Aquaculture* 136, 313–321.
- Uye S.-I. and Ichino S.** (1995) Seasonal variations in abundance, size composition, biomass and production rate of *Oikopleura dioica* (Fol) (Tunicata: Appendicularia) in a temperate eutrophic inlet. *Journal of Experimental Marine Biology and Ecology* 189, 1–11.
- Valletín R.** (1907) The fauna of St Ives Bay. *Journal of the Royal Institution of Cornwall* LIII, 29.
- Verwey J.** (1967) Annual report of the Netherlands Institute for Sea Research for the year 1964. *Netherlands Journal of Sea Research* 3, 464–487.
- Williams G.** (1954) The fauna of Strangford Lough and neighbouring coasts. *Proceedings of the Royal Irish Academy* 56, 29–133.
- and
- Zaret T.M. and Kerfoot W.C.** (1975) Fish predation on *Bosmina longirostris*: body-size selection versus visibility selection. *Ecology* 56, 232–237.

#### Correspondence should be addressed to:

M.K.S. Lilley  
 Institute of Environmental Sustainability, Swansea University  
 Singleton Park, Swansea, SA2 8PP, UK  
 email: 385124@swansea.ac.uk